

GUÍA DE ESTUDIO: ARTÍCULOS

MAMA: In situ

- Godwin A. Post-operative radiotherapy for ductal carcinoma in situ of the breast (Review). **Cochrane Database Syst Rev.** 2013 Nov 21;(11):CD000563
- McCormick B. **RTOG 9804:** a prospective randomized trial for good-risk ductal carcinoma in situ comparing radiotherapy with observation. **J Clin Oncol.** 2015 Mar 1;33(7):709-715.

MAMA: Temprano

- **EBCTCG.** Effect of radiotherapy after breast-conserving surgery on 10-year recurrence and 15-year breast cancer death: meta-analysis of individual patient data for 10 801 women in 17 randomized trials. **Lancet.** 2011 Nov 12; 378(9804):1707–1716.
- Polo A. Risk factors and state-of-the-art indications for boost irradiation in invasive breast carcinoma. **Brachytherapy** 2017.
- Haviland JS. The UK Standardisation of Breast Radiotherapy (**START**) trials of radiotherapy hypofractionation for treatment of early breast cancer: 10-year follow-up results of two randomised controlled trials. **Lancet Oncol.** 2013 Oct;14(11):1086-1094.
- START Trialists Group. The UK Standardisation of Breast Radiotherapy (**START**) Trial **A** of radiotherapy hypofractionation for treatment of early breast cancer: a randomised trial. **Lancet Oncol** 2008; 9: 331–341.
- START Trialists Group. The UK Standardisation of Breast Radiotherapy (**START**) Trial **B** of radiotherapy hypofractionation for treatment of early breast cancer: a randomised trial. **Lancet** 2008; 371: 1098–1107.
- Smith BD. Radiation therapy for the whole breast: Executive summary of an American Society for Radiation Oncology (**ASTRO**) evidence-based guideline. **Practical Radiation Oncology** 2018;8:145-152.
- Veronesi U. Intraoperative radiotherapy versus external radiotherapy for early breast cancer (**ELIOT**): a randomised controlled equivalence trial. **Lancet Oncol** 2013;14:129-1277.
- Correa C. Accelerated Partial Breast Irradiation: Executive summary for the update of an **ASTRO** Evidence-Based Consensus Statement. **Pract Radiat Oncol** 2017;7(2):73-79.
- Kunkler IH. Breast-conserving surgery with or without irradiation in women aged 65 years or older with early breast cancer (**PRIME II**): a randomised controlled trial. **Lancet Oncol** 2015;3(16):266-273

MAMA: Localmente avanzado

- Nielsen HM. Study of Failure Pattern Among High-Risk Breast Cancer Patients With or Without Postmastectomy Radiotherapy in Addition to Adjuvant Systemic

Therapy: Long-Term Results From the Danish Breast Cancer Cooperative Group DBCG 82 b and c Randomized Studies. *J Clin Oncol* 2006;15(24): 2268-2274

- Overgaard M. Postoperative radiotherapy in high-risk premenopausal women with breast cancer who receive adjuvant chemotherapy. *NEJM* 1997;14(337):949-955
- Overgaard M. Postoperative radiotherapy in high-risk postmenopausal breast cancer patients given adjuvant tamoxifen: Danish Breast Cancer Cooperative Group DBCG 82c randomised trial. *Lancet* 1999;353:1641-1648
- Recht A. Postmastectomy Radiotherapy: An American Society of Clinical Oncology, American Society for Radiation Oncology, and Society of Surgical Oncology Focused Guideline Update. *JCO* 2016;36(34):4431-4442.

GINECOLOGÍA: Cancer de cérvix

- Rotman M. A phase III randomized trial of postoperative pelvic irradiation in stage IB cervical carcinoma with poor prognostic features: follow-up of a Gynecologic Oncology Group study. **GOG-92**. *IJROBP* 2006;1(65):169-176
- Peters WA. Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix. **GOG 109**. *JCO* 2000;18:1606-1613
- Landoni F. Randomized study between radical surgery and radiotherapy for the treatment of stage IB-IIA cervical cancer: 20-year update. *J Gynecol Oncol*. 2017 May;28(3):e34
- Girinsky T. Overall treatment time in advanced cervical carcinomas: a critical parameter in treatment outcome. *IJROBP* 1993;27:1051-1056
- Eifel PJ. Pelvic irradiation with concurrent chemotherapy versus pelvic and para-aortic irradiation for high-risk cervical cancer: an update of radiation therapy oncology group trial (**RTOG 90-01**). *J Clin Oncol* 2004; 22:872-880.
- Rose PG. Long-term follow-up of a randomized trial comparing concurrent single agent cisplatin, cisplatin-based combination chemotherapy, or hydroxyurea during pelvic irradiation for locally advanced cervical cancer: a Gynecologic Oncology Group Study. *J Clin Oncol* 25:2804-2810.

GINECOLOGÍA: Cáncer de endometrio

- Keys HM. A phase III trial of surgery with or without adjunctive external pelvic radiation therapy in intermediate risk endometrial adenocarcinoma: a Gynecologic Oncology Group study. *Gynecologic Oncology* 2004;92:744-751.
- Creutzberg CL. Surgery and postoperative radiotherapy versus surgery alone for patients with stage-1 endometrial carcinoma: multicentre randomised trial. *Lancet* 2000; 355: 1404-1411.
- Nout RA. Vaginal brachytherapy versus pelvic external beam radiotherapy for patients with endometrial cancer of high-intermediate risk (PORTEC-2): an open-label, non-inferiority, randomised trial. *Lancet* 2010; 375: 816-823.

- de Boer SM. Adjuvant chemoradiotherapy versus radiotherapy alone for women with high-risk endometrial cancer (PORTEC-3): final results of an international, open-label, multicentre, randomised, phase 3 trial. *Lancet* 2018;3(19):295-309.

GINECOLOGÍA: Cáncer de vulva

- Sciacero P. The role of radiation therapy in vulvar cancer: review of the current literature. *Tumori* 2016;103(5):422-429.
- Chapman BV. Adjuvant radiation therapy for margin-positive vulvar squamous cell carcinoma: defining the ideal dose-response using the national cancer data base. *Int J Radiation Oncol Biol Phys* 2017; 97(1): 107-117.

NEUMOLOGÍA: Cáncer de pulmón de células no pequeñas

- Timmerman R. Stereotactic body radiation therapy for inoperable early stage lung cancer. *JAMA* 2010;303(11):1070-1076.
- Nyman J. SPACE – A randomized study of SBRT vs conventional fractionated radiotherapy in medically inoperable stage I NSCLC. *Radiotherapy and Oncology* 2016;121:1-8.
- Chang JY. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Oncol* 2015; 16: 630–637.
- Videtic GMM. Stereotactic body radiation therapy for early stage non-small cell lung cancer: Executive Summary of an ASTRO Evidence-Based Guideline. *Practical Radiation Oncology* 2017;7:295-301.
- PORT Meta-analysis Trialists Group. Postoperative radiotherapy for non-small cell lung cancer (Review). *Cochrane Database Syst Rev* 2005 Apr 18;(2):CD002142
- Burdett S. PORT Meta-analysis Trialists Group. Postoperative radiotherapy for non-small cell lung cancer (Review). *Cochrane Database Syst Rev* 2016;9:CD002142
- Pless M. Induction chemoradiation in stage IIIA/N2 non-small-cell lung cancer: a phase 3 randomised trial. *Lancet* 2015; 386: 1049–1056.
- Aupérin A. Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. *J Clin Oncol* 2010;28:2181-2190.
- Bradley JD. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (**RTOG 0617**): a randomised, two-by-two factorial phase 3 study. *Lancet Oncol* 2015;16(2):187-199.
- Moeller B. Palliative thoracic radiation therapy for non-small cell lung cancer: 2018 Update of an American Society for Radiation Oncology (ASTRO) Evidence-Based Guideline. *Pract Radiat Oncol* 2018;8(4):245-250.

NEUMOLOGÍA: Cáncer de pulmón de células pequeñas

- Pignon JP. A Meta-Analysis of Thoracic Radiotherapy for Small-Cell Lung Cancer. *N Engl J Med* 1992; 327:1618-1624.

- Turrisi AT. Twice-Daily Compared with Once-Daily Thoracic Radiotherapy in Limited Small-Cell Lung Cancer Treated Concurrently with Cisplatin and Etoposide. *N Engl J Med* 1999; 340:265-271.
- Aupérin A. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. Prophylactic Cranial Irradiation Overview Collaborative Group. *N Engl J Med* 1999;341(7):476-484.
- Jeremic B. Role of Radiation Therapy in the Combined-Modality Treatment of Patients With Extensive Disease Small-Cell Lung Cancer: A Randomized Study. *J Clin Oncol* 1999;17:2092-2099.
- Slotman BJ. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: a phase 3 randomised controlled trial. *Lancet* 2015;385:36–42.
- Takahashi T. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2017;18(5):663-671.

UROLOGÍA: Cáncer de próstata

- Rahul D. Contemporary Update of a Multi-Institutional Predictive Nomogram for Salvage Radiotherapy After Radical Prostatectomy. *J Clin Oncol* 34:3648-3654.
- Bolla M. Postoperative radiotherapy after radical prostatectomy for high-risk prostate cancer: long-term results of a randomized controlled trial (EORTC trial 22911). *Lancet* 2012; 380: 2018–2027.
- Hamdy FC. 10-Year Outcomes after Monitoring, Surgery, or Radiotherapy for Localized Prostate Cancer. *N Engl J Med* 2016;375:1415-1424.
- Barocas DA. Association Between Radiation Therapy, Surgery, or Observation for Localized Prostate Cancer and Patient-Reported Outcomes After 3 Years. *JAMA*. 2017;317(11):1126-1140.
- Catton CN. Randomized Trial of a Hypofractionated Radiation Regimen for the Treatment of Localized Prostate Cancer. *J Clin Oncol* 2017;35:1884-1890
- Morgan SC. Hypofractionated Radiation Therapy for Localized Prostate Cancer: Executive Summary of an ASTRO, ASCO, and AUA Evidence-Based Guideline. *Practical Radiation Oncology* 2018;8:354-360.
- Jones CU. Radiotherapy and short-term androgen deprivation for localized prostate cancer. *N Engl J Med* 2011;365(2):107-118
- Dearnaley DP. Escalated-dose versus control-dose conformal radiotherapy for prostate cancer: long-term results from the MRC RT01 randomised controlled trial. *Lancet Oncol* 2014; 15: 464–473.
- Bolla M. External irradiation with or without long-term androgen suppression for prostate cancer with high metastatic risk: 10-year results of an EORTC randomised study. *Lancet Oncol* 2010; 11: 1066–1073.
- Parker CC. Radiotherapy to the primary tumour for newly diagnosed, metastatic prostate cancer (STAMPEDE): a randomised controlled phase 3 trial. *Lancet* 2018; 392: 2353–2366.

UROLOGÍA: Seminoma

- Jones WG. Randomized trial of 30 versus 20 Gy in the adjuvant treatment of stage I Testicular Seminoma: a report on Medical Research Council Trial TE18, European Organisation for the Research and Treatment of Cancer Trial 30942 (ISRCTN18525328). *J Clin Oncol* 2005;23(6):1200-1208.
- Oliver RTD. Radiotherapy versus single-dose carboplatin in adjuvant treatment of stage I seminoma: a randomised trial. *Lancet* 2005; 366: 293–300.
- Oliver RTD. Randomized Trial of Carboplatin Versus Radiotherapy for Stage I Seminoma: Mature Results on Relapse and Contralateral Testis Cancer Rates in MRC TE19/EORTC 30982 Study (ISRCTN27163214). *J Clin Oncol* 2011;29:957-962.
- Hallemeier CL. Excellent long-term disease control with modern radiotherapy techniques for stage I testicular seminoma—The Mayo Clinic experience. *Urologic Oncology: Seminars and Original Investigations* 2014;32:24.e1–24.e6.
- Patel HD. Radiotherapy for stage I and II testicular seminomas: Secondary malignancies and survival. *Urol Oncol* 2017;35(10):606.e1-606.e7.

UROLOGÍA: Cáncer de vejiga

- James ND. Radiotherapy with or without Chemotherapy in Muscle-Invasive Bladder Cancer. *N Engl J Med* 2012;366:1477-1488.

CABEZA Y CUELLO: Posoperados

- Cooper JS. Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *N Engl J Med* 2004;350:1937-1944.
- Cooper JS. Long-term follow-up of the RTOG 9501/Intergroup phase III Trial: Postoperative concurrent radiation therapy and chemotherapy in high-risk squamous cell carcinoma of the head and neck. *Int J Radiation Oncol Biol Phys* 2012; 84(5):1198-1205.
- Bernier J. Postoperative irradiation with or without concomitant chemotherapy for locally advanced head and neck cancer. *N Engl J Med* 2004;350:1945-1952.
- Bernier J. Defining risk levels in locally advanced head and neck cancers: a comparative analysis of concurrent postoperative radiation plus chemotherapy trials of the EORTC (#22931) and RTOG (#9501). *Head Neck*. 2005;27(10):843-850.

CABEZA Y CUELLO: Localmente avanzado

- Pignon JP. Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): An update on 93 randomised trials and 17,346 patients. *Radiotherapy and Oncology* 2009;92:4–14.
- Beittler JJ. Final results of local-regional control and late toxicity of RTOG 9003: A randomized trial of altered fractionation radiation for locally advanced head and neck cancer. *Int J Radiation Oncol Biol Phys* 2014; 89(1): 13-20

CABEZA Y CUELLO: Cavidad oral y orofaringe

- Sher DJ. Radiation therapy for oropharyngeal squamous cell carcinoma: Executive summary of an ASTRO Evidence-Based Clinical Practice Guideline. *Practical Radiation Oncology*. 2017;7:246–253.

CABEZA Y CUELLO: Hipofaringe

- Lefebvre JL. Laryngeal preservation with induction chemotherapy for hypopharyngeal squamous cell carcinoma: 10-year results of EORTC trial 24891. *Annals of Oncology* 2012;23:2708–2714.

CABEZA Y CUELLO: Laringe

- Eskiizmir G. Risk factors for radiation failure in early-stage glottic carcinoma: A systematic review and meta-analysis. *Oral Oncology* 62 (2016) 90–100.
- Yamazaki H. Radiotherapy for early glottic carcinoma (T1N0M0): Results of prospective randomized study of radiation fraction size and overall treatment time. *Int J Radiation Oncology Biol Phys* 2006; 64(1):77–82.
- Trott A. Randomized trial of hyperfractionation versus conventional fractionation in T2 squamous cell carcinoma of the vocal cord (RTOG 9512). *Int J Radiation Oncol Biol Phys* 2014;89(5):958-963.
- **VETERANS.** Wolf GT. Induction chemotherapy plus radiation compared with surgery plus radiation in patients with advanced laryngeal cancer. *N Engl J Med* 1991;324(24):1685-1690.
- Forastiere AA. Long-term results of **RTOG 91-11**: A comparison of three nonsurgical treatment strategies to preserve the larynx in patients with locally advanced larynx cancer. *J Clin Oncol* 2012;31:845-852.

CABEZA Y CUELLO: Nasofaringe

- Al-Sarraf M. Chemoradiotherapy versus radiotherapy in patients with advanced nasopharyngeal cancer: Phase III Randomized **Intergroup Study 0099**. *J Clin Oncol* 1998;16:1310-1317.
- Wee J. Randomized trial of radiotherapy versus concurrent chemoradiotherapy followed by adjuvant chemotherapy in patients with American Joint Committee on Cancer/International Union against cancer stage III and IV nasopharyngeal cancer of the endemic variety. **SQNP01**. *J Clin Oncol* 2005;23(27):6730-6738.
- Lee AWM. A multicenter, phase 3, randomized trial of concurrent chemoradiotherapy plus adjuvant chemotherapy versus radiotherapy alone in patients with regionally advanced nasopharyngeal carcinoma: 10-year outcomes for efficacy and toxicity. **NPC-9901**. *Cancer* 2017;123(21):4147-4157.
- Lee N. Intensity-modulated radiation therapy with or without chemotherapy for nasopharyngeal carcinoma: **Radiation Therapy Oncology Group phase II trial 0225**. *J Clin Oncol* 2009;27:3684-3690.

CABEZA Y CUELLO: Glándulas salivales

- Terhaard CH. Postoperative and primary radiotherapy for salivary gland carcinomas: indications, techniques, and results. *Int J Radiat Oncol Biol Phys* 2007;69(2 Suppl):S52-S55.
- Mahmood U. Adjuvant Radiation Therapy for High-Grade and/or Locally Advanced Major Salivary Gland Tumors. *Arch Otolaryngol Head Neck Surg* 2011;137(10):1025-1030.

CABEZA Y CUELLO: Tiroides

- Kiess AP. External-beam radiotherapy for differentiated thyroid cancer locoregional control: A statement of the American Head and Neck Society. *Head Neck*. 2016 Apr;38(4):493-498.
- Fussey JM. External beam radiotherapy in differentiated thyroid carcinoma: A systematic review. *Head Neck* 2016;38:E2297–E2305
- Mangoni M. External beam radiotherapy in thyroid carcinoma: clinical review and recommendations of the AIRO “Radioterapia Metabolica” Group. *Tumori* 2017; 103(2): 114-123.
- Kwon J. The prognostic impacts of postoperative radiotherapy in the patients with resected anaplastic thyroid carcinoma: A systematic review and meta-analysis. *European Journal of Cancer* 2016;59: 34-45

NEUROLOGÍA: Metástasis a sistema nervioso central

- Kocher M. Adjuvant whole-brain radiotherapy versus observation after radiosurgery or surgical resection of one to three cerebral metastases: results of the **EORTC 22952-26001** study. *J Clin Oncol*. 2011 Jan 10;29(2):134-141.
- Tsao MN. Radiotherapeutic and surgical management for newly diagnosed brain metastasis(es): **An American Society for Radiation Oncology evidence-based guideline**. *Pract Radiat Oncol*. 2012 Jul; 2(3): 210–225.
- Patil CG. Whole brain radiation therapy (WBRT) alone versus WBRT and radiosurgery for the treatment of brain metastases (Review). **Cochrane Database of Syst Rev** 2017;9:CD006121
- Lamba N. Stereotactic radiosurgery versus whole brain radiotherapy after intracranial metastasis resection: a systematic review and meta-analysis. *Radiation Oncology* 2017;12:106

NEUROLOGÍA: Gliomas alto grado

- Lecavalier-Barsoum M. Cochrane Database Syst Rev. 2014 May 15;(5):CD007104. Adjuvant treatment of anaplastic oligodendrogiomas and oligoastrocytomas.

NEUROLOGÍA: Glioblastoma

- Stupp R. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009;10:459–466

- Malmström A. Temozolomide versus standard 6-week radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 2012; 13: 916–926.
- Perry J. Short-course radiation plus temozolomide in elderly patients with glioblastoma. *N Engl J Med* 2017;376:1027-1037.

NEUROLOGÍA: Gliomas bajo grado

- Baumert BG. Temozolomide chemotherapy versus radiotherapy in high-risk low-grade glioma (**EORTC 22033-26033**): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol* 2016; 17: 1521–1532.
- Buckner JC. Radiation plus Procarbazine, CCNU, and Vincristine in Low-Grade Glioma. **RTOG 9802**. *N Engl J Med* 2016;374:1344-1355.
- van den Bent MJ. Long-term efficacy of early versus delayed radiotherapy for low-grade astrocytoma and oligodendrogloma in adults: the **EORTC 22845** randomised trial. *Lancet* 2005; 366: 985–990.

NEUROLOGÍA: Medulloblastoma

- Packer RJ. Phase III study of craniospinal radiation therapy followed by adjuvant chemotherapy for newly diagnosed average-risk medulloblastoma. *J Clin Oncol*. 2006;24(25):4202-4208.
- Ashley DM. Induction chemotherapy and conformal radiation therapy for very young children with nonmetastatic medulloblastoma: Children's Oncology Group study P9934. *J Clin Oncol*. 2012 Sep 10;30(26):3181-3186.

NEUROLOGÍA: Cordinoma

- Sahgal A. Image-guided, intensity-modulated radiation therapy (IG-IMRT) for skull base chordoma and chondrosarcoma: preliminary outcomes. *Neuro-Oncology* 2015;17(6):889–894.

GASTROENTEROLOGÍA: Esófago

- Cooper JS. Chemoradiotherapy of locally advanced esophageal cancer: long-term follow-up of a prospective randomized trial (**RTOG 85-01**). *Radiation Therapy Oncology Group*. *JAMA* 1999;281(17):1623-1627.
- Zhu LL. A Meta-Analysis of Concurrent Chemoradiotherapy for Advanced Esophageal Cancer. *PLoS ONE* 10(6): e0128616.
- Minsky BD. **INT 0123** (Radiation Therapy Oncology Group 94-05) Phase III trial of combined-modality therapy for esophageal cancer: high-dose versus standard-dose radiation therapy. *J Clin Oncol* 2002;5(20):1167-1174.
- Shapiro J. Neoadjuvant chemoradiotherapy plus surgery versus surgery alone for oesophageal or junctional cancer (**CROSS**): long-term results of a randomised controlled trial. *Lancet* 2015; 16(9):1090-1098.

GASTROENTEROLOGÍA: Estómago

- MacDonald JS. Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med* 2001;345:725-730.
- Smalley SR. Updated analysis of SWOG-directed Intergroup Study 0116: A phase III trial of adjuvant radiochemotherapy versus observation after curative gastric cancer resection. *J Clin Oncol*. 2012 Jul 1;30(19):2327-2333.
- Lee J. Phase III trial comparing capecitabine plus cisplatin versus capecitabine plus cisplatin with concurrent capecitabine radiotherapy in completely resected gastric cancer with D2 lymph node dissection: The ARTIST Trial. *J Clin Oncol* 2011;30:268-273.
- Leong T. TOPGEAR: A randomized, Phase III trial of perioperative ECF chemotherapy with or without preoperative chemoradiation for resectable gastric cancer: interim results from an international, intergroup trial of the AGITG, TROG, EORTC and CCTG. *Ann Surg Oncol*. 2017;24(8):2252-2258.

GASTROENTEROLOGÍA: Recto

- Sauer R. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 2004;351:1731-1740.
- Sauer R. Preoperative Versus Postoperative Chemoradiotherapy for Locally Advanced Rectal Cancer: Results of the German CAO/ARO/AIO-94 Randomized Phase III Trial After a Median Follow-Up of 11 Years. *J Clin Oncol* 2012;30:1926-1933.
- Erlandsson J. Optimal fractionation of preoperative radiotherapy and timing to surgery for rectal cancer (Stockholm III): a multicentre, randomised, non-blinded, phase 3, non-inferiority trial. *Lancet Oncol*. 2017;18(3):336-346.

GASTROENTEROLOGÍA: Canal anal

- Nigro ND. An evaluation of combined therapy for squamous cell cancer of the anal canal. *Diseases of the colon & rectum*. 1984;12(27):763-766
- Northover J. Chemoradiation for the treatment of epidermoid anal cancer: 13-year follow-up of the first randomised UKCCR Anal Cancer Trial (**ACT I**). *British Journal of Cancer* 2010;102:1123-1128
- Glynne-Jones R. Best time to assess complete clinical response after chemoradiotherapy in squamous cell carcinoma of the anus (**ACT II**): a post-hoc analysis of randomised controlled phase 3 trial. *Lancet Oncol* 2017;18(3):347-356

GASTROENTEROLOGÍA: Cáncer de páncreas

- Tang K. Neoadjuvant therapy for patients with borderline resectable pancreatic cancer: A systematic review and meta-analysis of response and resection percentages. *Pancreatology* 2016;16(1):28-37.
- Neoptolemos JP. A Randomized Trial of Chemoradiotherapy and Chemotherapy after Resection of Pancreatic Cancer. *N Engl J Med* 2004;350:1200-1210.

- van Tienhoven G. Preoperative chemoradiotherapy versus immediate surgery for resectable and borderline resectable pancreatic cancer (PREOPANC-1): A randomized, controlled, multicenter phase III trial. *J Clin Oncol* 36(18):LBA4002-LBA4002

GASTROENTEROLOGÍA: Hígado y vía biliar

- Scorsetti M. Stereotactic body radiation therapy for liver metastases. *J Gastrointest Oncol* 2014;5(3):190-197
- Murray LJ. Advances in Stereotactic Body Radiation Therapy for Hepatocellular Carcinoma. *Semin Radiat Oncol* 2017;27:247-255.
- Horgan AM. Adjuvant therapy in the treatment of biliary tract cancer: A systematic review and meta-analysis. *J Clin Oncol* 2012;30:1934-1940.

PIEL Y TEJIDOS BLANDOS: Melanoma

- Agrawal S. The benefits of adjuvant radiation therapy after therapeutic lymphadenectomy for clinically advanced, high-risk, lymph node-metastatic melanoma. *Cancer* 2009;115:5836-5844.
- Burmeister BH. Adjuvant radiotherapy versus observation alone for patients at risk of lymph-node field relapse after therapeutic lymphadenectomy for melanoma: a randomised trial. *Lancet Oncol* 2012; 13: 589-597.
- Henderson MA. Adjuvant lymph-node field radiotherapy versus observation only in patients with melanoma at high risk of further lymph-node field relapse after lymphadenectomy (ANZMTG 01.02/TROG 02.01): 6-year follow-up of a phase 3, randomised controlled trial. *Lancet Oncol* 2015; 16: 1049-1060.

PIEL Y TEJIDOS BLANDOS: Sarcoma de tejido blando

- O'Sullivan Brian. Preoperative versus postoperative radiotherapy in soft-tissue sarcoma of the limbs: a randomised trial. *Lancet* 2002, 359: 2235-2241.
- Albertsmeier M. External beam radiation therapy for resectable soft tissue sarcoma: A systematic review and meta-analysis. *Ann Surg Oncol* 2018;25(3):754-767.

PIEL Y TEJIDOS BLANDOS: Osteosarcoma

- Sole CV. Adjuvant radiation therapy in resected high-grade localized skeletal osteosarcomas treated with neoadjuvant chemotherapy: Long-term outcomes. *Radiother Oncol* 2016;119(1):30-34.

HEMATOLOGÍA: Linfoma de Hodgkin

- Engert A. Two cycles of doxorubicin, bleomycin, vinblastine, and dacarbazine plus extended-field radiotherapy is superior to radiotherapy alone in early favorable Hodgkin's Lymphoma: final results of the GHSG HD7 Trial. *J Clin Oncol* 2007;25:3495-3502.

- Engert A. Reduced treatment intensity in patients with early-stage hodgkin's lymphoma. N Engl J Med 2010;363:640-652.
- Eich HT. Intensified chemotherapy and dose-reduced involved-field radiotherapy in patients with early unfavorable hodgkin's lymphoma: Final analysis of the German Hodgkin Study Group HD11 Trial. J Clin Oncol 2010;28:4199-4206
- Borchmann P. Eight cycles of escalated-dose BEACOPP compared with four cycles of escalated-dose BEACOPP followed by four cycles of baseline-dose BEACOPP with or without radiotherapy in patients with advanced-stage hodgkin's lymphoma: final analysis of the HD12 Trial of the German Hodgkin Study Group. J Clin Oncol 2011;29:4234-4242.
- Raemaekers JM. Omitting radiotherapy in early positron emission tomography-negative stage I/II Hodgkin lymphoma is associated with an increased risk of early relapse: Clinical results of the preplanned interim analysis of the randomized EORTC/LYSA/FIL H10 trial. J Clin Oncol. 2014 Apr 20;32(12):1188-1194.

HEMATOLOGÍA: Linfoma no Hodgkin

- Lowry L. Reduced dose radiotherapy for local control in non-Hodgkin lymphoma: A randomised phase III trial. Radiotherapy and Oncology 2011;100:86–92.
- Hoskin PJ. 4 Gy versus 24 Gy radiotherapy for patients with indolent lymphoma (FORT): a randomised phase 3 non-inferiority trial. Lancet Oncol 2014;15(4):457-463.
- Horning SJ. Chemotherapy with or without radiotherapy in limited-stage diffuse aggressive non-Hodgkin's lymphoma: Eastern Cooperative Oncology Group Study 1484. J Clin Oncol 2004;22:3032-3038.
- Miller TP. Chemotherapy alone compared with chemotherapy plus radiotherapy for localized intermediate- and high-grade non-hodgkin's lymphoma. N Engl J Med 1998;339:21-26

PEDIATRÍA: Rabdomiosarcoma

- Raney RB. The Intergroup Rhabdomyosarcoma Study Group (IRS): major lessons from the IRS-I through IRS-IV studies as background for the current IRS-V treatment protocols. Sarcoma 2001;5:9-15

PEDIATRÍA: Sarcoma de Ewing

- Foulon S. Can postoperative radiotherapy be omitted in localised standard-risk Ewing sarcoma? An observational study of the Euro-E.W.I.N.G group. European Journal of Cancer 2016; 61 :128-136

OTROS: metástasis óseas

- Lutz S. Palliative radiation therapy for bone metastases: Update of an ASTRO Evidence-Based Guideline. Practical Radiation Oncology 2017;7:4-12.

OTROS: compresión medular

- Patchell RA. Direct decompressive surgical resection in the treatment of spinal cord compression caused by metastatic cancer: a randomised trial. Lancet 2005;366:643-648
- Chen B. Comparison of the therapeutic efficacy of surgery with or without adjuvant radiotherapy versus radiotherapy alone for metastatic spinal cord compression: A meta-analysis. World Neurosurg 2015;83(6):1066-1073.
- Rades D. Evaluation of Five Radiation Schedules and Prognostic Factors for Metastatic Spinal Cord Compression. J Clin Oncol 2005;23:3366-3375.
- Rades D. Radiotherapy With 4 Gy x 5 Versus 3 Gy x 10 for Metastatic Epidural Spinal Cord Compression: Final Results of the SCORE-2 Trial (ARO 2009/01). J Clin Oncol 2016; 34.